



U.S. ARMY MEDICAL RESEARCH INSTITUTE OF CHEMICAL DEFENSE

USAMRICD-TR-96-03

Autoradiographic Localization of [¹²⁵I]-Ricin in Lungs and Trachea of Mice Following an Aerosol Inhalation Exposure

Jeffrey A. Doeblner
Robert B. Moeller
Norman D. Wiltshire
Thomas W. Mayer
James E. Estep
Richard K. Traub
Clarence A. Broomfield
Craig A. Calamario
William L. Thompson
M. Louise Pitt

DTIC QUALITY INSPECTED 4

June 1996

19960813 107

Approved for public release; distribution unlimited

U.S. Army Medical Research
Institute of Chemical Defense
Aberdeen Proving Ground, MD 21010-5425

DISPOSITION INSTRUCTIONS:

Destroy this report when no longer needed. Do not return to the originator.

DISCLAIMERS:

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

In conducting the work described in this report, the investigators adhered to the "Guide for the Care and Use of Laboratory Animals," NIH Publication 86-23, revised in 1985.

The use of trade names does not constitute an official endorsement or approval of the use of such commercial hardware or software. This document may not be cited for purposes of advertisement.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE June 1996	3. REPORT TYPE AND DATES COVERED Technical Report	
4. TITLE AND SUBTITLE Autoradiographic Localization of [¹²⁵ I]-Ricin in Lungs and Trachea of Mice Following an Aerosol Inhalation Exposure		5. FUNDING NUMBERS 61102A 3M161102BS12 AA	
6. AUTHOR(S) Doeblner, JA, Moeller, RB, Wiltshire, ND, Mayer, TW, Estep, JE, Traub, RK, Broomfield, CA, Calamaio, CA, Thompson, WL, Pitt, ML			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Commander U.S. Army Medical Research Institute of Chemical Defense ATTN: MCMR-UV-PN 3100 Ricketts Point Road Aberdeen Proving Ground, MD 21010-5425		8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Commander U.S. Army Medical Research Institute of Chemical Defense ATTN: MCMR-UV-RC 3100 Ricketts Point Road Aberdeen Proving Ground, MD 21010-5425		10. SPONSORING/MONITORING AGENCY REPORT NUMBER USAMRICD-TR-96-03	
11. SUPPLEMENTARY NOTES			
12a. DISTRIBUTION AVAILABILITY STATEMENT Approved for public release; distribution unlimited		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Studies were conducted to examine the light microscopic distribution of [¹²⁵ I]-ricin within the lungs and corresponding lung pathology in mice following single, nose-only, aerosol inhalation exposures. Groups of 4-8 mice were euthanatized 15 minutes and 1, 4, 6, 8, 12, and 24 hours post-exposure. At 15 minutes post-exposure, abundant radiolabel was present within alveolar spaces. At all later times, however, [¹²⁵ I]-ricin was observed primarily within the lumina of bronchioles and larger passageways. Cellular targets for ricin within the lungs could not be identified using the techniques employed; it is therefore recommended that ultrastructural autoradiography be used to determine cellular ricin binding targets. The primary pathological abnormality detected was multifocal epithelial necrosis of the secondary bronchioles.			
14. SUBJECT TERMS Ricin, Autoradiography, Lungs, Aerosol Inhalation		15. NUMBER OF PAGES 10	
		16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UNLIMITED

Acknowledgments

The authors wish to thank H. Shafer for excellent technical support.

Introduction

The inhalation of aerosolized ricin is believed to represent a potentially significant hazard to military personnel (Cope et al., 1946; Quinchon and Levy, 1962; Korosuo, 1971). The primary biochemical lesion produced by ricin is inactivation of protein synthesis (Olsnes et al., 1974). However, the precise sites of toxin action remain unknown. Recent studies conducted in our laboratories (Doebler et al., 1995) have suggested that the lungs may represent the only pathophysiologically important site of [125 I]-ricin accumulation following an aerosol exposure. This hypothesis is consistent with previous results indicating that histopathological abnormalities are essentially restricted to the lungs following an aerosol exposure to ricin (Griffiths et al., 1993).

The current investigation was conducted to determine the microscopic distribution of [125 I]-ricin in mice following aerosol inhalation exposures and to relate ricin binding to pathological findings.

Materials and Methods

Ricin (Vector Laboratories, Burlingame, CA) was iodinated with carrier-free [125 I] (Amersham, Arlington Heights, IL) using the lactoperoxidase method (Marchalonis, 1969). Extensive studies were conducted previously to validate the use of [125 I]-ricin as a probe for toxin localization (Doebler et al., 1995). The toxicity of [125 I]-ricin was confirmed both *in vitro* (using inhibition of protein synthesis and inhibition of cellular proliferation as endpoints) and *in vivo* (using a two-dosage regimen with survival as endpoint). In addition, the degradation of [125 I]-ricin in mouse tissues was found to be substantial (69-74% of radiolabel in liver and spleen was precipitated in 10% trichloroacetic acid at 60 min post-injection), but not so severe as to preclude the use of [125 I]-ricin as a probe for toxin localization (Doebler et al., 1995).

Twenty-eight male CD-1 mice (weighing 18-22 grams) were obtained from Harlan Sprague-Dawley Inc. (Detrick, MD). Upon arrival, mice were quarantined and observed for evidence of disease for 5 days before use. Mice were housed in polycarbonate shoebox cages in groups of 10 or less per cage. They were provided commercial certified mouse ration and tap water ad libitum. Animal holding rooms were maintained at 21 ± 2 °C on a 12-hour light/dark cycle. All mice received [125 I]-ricin (specific activity 1.72×10^9 cpm/mg) by nose-only aerosol inhalation. Groups of 4 mice were euthanatized (as described below) at 15 minutes, 1, 4, 8, 12 and 24 hours after [125 I]-ricin exposure; an additional group of 4 mice was euthanatized at 1 hour post-exposure (total n for 1 hour = 8). Aerosols of ricin (13 mg/minute/meter³) were generated by a Collison nebulizer operated at 26 pounds per square inch (psi) of air pressure, giving 7.5 liters/minute with mass median aerodynamic diameter (MMAD) of 1.06 μ m (geometric standard deviation = 1.203) as measured by an Aerodynamic Particle Sizer, APS3000 (TSI Inc., St. Paul, MN). Total flow rate through the non-rebreathing system was adjusted to 12 liters/minute with a secondary air flow of 4.5 liters/minute. Aerosol sampling was conducted continuously for the duration of the 10-minute exposure using an all glass impinger (AGI) with a sonic velocity of 6.0 liters/minute. The ricin exposure level was sublethal (Estep, unpublished data) and did not produce overt signs of toxicity in any of the exposed animals.

Mice were euthanatized by pentobarbital overdose (6.5 mg/mouse, ip), and blood was collected (until heart stoppage) by snipping the tail. Lungs were removed in toto, frozen immediately in a dry ice-ethanol slurry and stored on dry ice. Entire lungs were mounted and sectioned (at 10 μ m) on a cryostatic microtome. Approximately every 10th or 15th section was saved onto a glass slide; this frequency was determined by the presence of large intrapulmonary passageways in the immediate vicinity of the section.

Preliminary studies were conducted to determine whether sufficient resolution could be obtained using X-ray film for the autoradiography of [125 I]-ricin administered by aerosol inhalation in the lungs of mice. Dupont (Wilmington, DE) Chronex and Ultra-Vision G films were placed in direct apposition to cryostatic sections of lung obtained from exposed mice. However, following development of the autoradiograms, it was obvious that the resolution provided by either film was not sufficient to localize the [125 I] label into various compartments of the lung. Thus, an autoradiographic emulsion (Kodak NTB-3 Nuclear Track Emulsion, Rochester, NY) was used in subsequent studies for the localization of [125 I]-ricin. Tissues on slides were covered with NTB-3 emulsion, placed vertically to allow even drying, and stored in the dark for 28 days. Routine developing procedures (using Kodak D-19 developer) were employed at the end of the exposure period. Sections were then counterstained with hematoxylin-eosin for microscopic examination.

Results and Discussion

Despite the markedly enhanced resolution provided by the emulsion relative to the films, specific cellular compartments accumulating [125 I]-ricin within the lung could not be identified with any degree of certainty. However, the label could be readily identified in specific intrapulmonary air passageways. This distribution is briefly described below.

The distribution of [125 I]-ricin within the lungs differed markedly among groups of mice sacrificed at 15 minutes and those groups sacrificed at all later times. In the 15-minute post-exposure group, abundant radiolabeled material was detected within the alveolar spaces and in the lumina of bronchioles, bronchi and the trachea. At later time periods (1-24 hours post-exposure), [125 I]-ricin was essentially restricted to the lumina of bronchioles and larger respiratory passageways. The disappearance of radiolabel from the alveolar spaces after the early post-exposure interval may signify the actual absorption of [125 I]-ricin into the alveoli.

No pathologic abnormalities could be detected in the lungs of mice sacrificed prior to 24 hours post-exposure. However, bronchiolar lesions were detected at 24 hours post-exposure. These bronchiolar lesions consisted of sites of minimal multifocal epithelial necrosis (observed primarily in secondary bronchioles) and were characterized by condensation of cytoplasm, pyknosis and disorganization of the epithelium. Some sloughing of the epithelium was also present. Occasional neutrophils were observed transmigrating the mucosa and submucosa.

The major objective of these studies was to determine the cellular distribution of [125 I]-ricin binding and corresponding histopathological abnormalities within the lungs of mice following nose-only aerosol inhalation exposures. However, the procedures employed were not sensitive enough to localize toxin or toxin-induced damage to specific cell types within the lungs. We believe that such information could be readily obtained using electron microscopic

autoradiography combined with routine morphological analyses. With regard to toxin localization, we have found that light microscopic autoradiography, performed with either film or emulsions, did not provide sufficient resolution to identify specific cell types. Since previous attempts to localize ricin using immunocytochemical techniques have also not been successful (D.R. Franz, personal communication), it appears that ultrastructural autoradiographic procedures, such as those described by Salpeter and Bachmann (1972), are required to determine ricin localization within specific cell types. We were also unable to identify any specific respiratory epithelial components within the lung that may have been selectively damaged by aerosol-administered ricin. The relatively thick (10 μ m) frozen sections made detailed evaluation of respiratory epithelial subtypes difficult. It is likely that routine morphological processing procedures (i.e., formalin-fixed tissues and 5-7 μ m paraffin sections) can be used to obtain additional insight as to specific lung cells damaged by aerosolized ricin.

References

- Cope, A.C., Dec, J., Cannan, R.K., Renshaw, B and Moore, S. (1946) Ricin. In: Chemical Warfare Agents and Related Chemical Problems, U.S. Office of Science Research and Development, National Defense Research Committee, Washington, DC, pp. 179-203.
- Doebler, J.A., Wiltshire, N.D., Mayer, T.W., Estep, J.E., Moeller, R.B., Traub, R.K., Broomfield, C.A., Calamaio, C.A., Thompson, W.L. and Pitt, M.L. (1995). The distribution of [¹²⁵I]-ricin in mice following aerosol inhalation exposure. *Toxicology* 98:137-149.
- Griffiths, G.D., Lindsay, C.D., Rice P. and Upshall, D.G. (1993) The toxicology of ricin and abrin toxins - studies on immunisation against abrin toxicity, in Proceedings of the 1993 Medical Defense Bioscience Review, U.S. Army Medical Research Institute of Chemical Defense, Aberdeen Proving Ground, MD, pp. 1407-1416. AD A275669.
- Korosuo, T., (1971) Review of Chemical Warfare. *Sothlaslaaketiet. Aik, L.*, 46(1):21-26.
- Marchalonis, J.J. (1969) An enzymic method for the trace iodination of immunoglobulins and other proteins. *Biochem. J.* 113:299-305.
- Olsnes, S., Refnes, K. and Pihl, A. (1974) Mechanism of action of the toxic lectins abrin and ricin. *Nature* 249:627-631.
- Quinchon, J. and Levy, R. (1962). Agents of Chemical Warfare: Properties and Possibilities for Manufacture. *Memorial des poudres* 44:157-173, 1962.
- Salpeter, M.M. and Bachmann, L. (1972). Autoradiography. In: Principles and Techniques of Electron Microscopy, Biological Applications. M.A. Hayat, ed. Van Nostrand-Reinhold Co., Inc., New York. pp. 221-278.

DISTRIBUTION LIST

Addresses	Copies	Addresses	Copies
DEFENSE TECHNICAL INFORMATION CENTER ATTN DTIC OCP 8725 JOHN J KINGMAN RD STE 0944 FT BELVOIR VA 22060-6218	2	DIRECTOR ARMED FORCES MEDICAL INTELLIGENCE CENTER FORT DETRICK MD 21702-5004	1
COMMANDER US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND FORT DETRICK MD 21702-5012	2	COMMANDER US ARMY INSTITUTE OF DENTAL RESEARCH BUILDING 40 WASHINGTON DC 20307-5300	1
HQDA DASG HCD WASHINGTON DC 20310	1	COMMANDER US ARMY INSTITUTE OF SURGICAL RESEARCH BUILDING 2653 FORT SAM HOUSTON TX 78234-6200	1
DIRECTOR WALTER REED ARMY INSTITUTE OF RESEARCH BUILDING 40 WASHINGTON DC 20307-5100	1	COMMANDANT ACADEMY OF HEALTH SCIENCES US ARMY ATTN HSHA CDC FORT SAM HOUSTON TX 78234-6100	1
COMMANDER US ARMY AEROMEDICAL RESEARCH LABORATORY ATTN SCIENTIFIC INFORMATION CENTER PO BOX 577 FORT RUCKER AL 36362-5000	1	COMMANDANT ACADEMY OF HEALTH SCIENCES US ARMY ATTN HSHA CDM FORT SAM HOUSTON TX 78234-6100	1
COMMANDER US ARMY MEDICAL RESEARCH INSTITUTE OF INFECTIOUS DISEASES BUILDING 1425 FORT DETRICK MD 21702-5011	1	Dr. JOSEPH OSTERMAN DIRECTOR ENVIRONMENTAL AND LIFE SCIENCES OFFICE OF THE DEPUTY DIRECTOR FOR RESEARCH AND ENGINEERING ROOM 3D129 WASHINGTON DC 20301-2300	1
COMMANDER US ARMY RESEARCH INSTITUTE OF ENVIRONMENTAL MEDICINE BUILDING 42 NATICK MA 01760-5007	1	COMMANDER US ARMY TRAINING AND DOCTRINE COMMAND ATTN ATMD FORT MONROE VA 23651	1
COMMANDANT US ARMY CHEMICAL SCHOOL ATTN ATZN CM C FORT MCCLELLAN AL 36205	1		

COMMANDER US ARMY NUCLEAR AND CHEMICAL AGENCY 7500 BACKLICK ROAD BUILDING 2073 SPRINGFIELD VA 22150-3198	1	AFOSR/NL BUILDING RM A217 BOLLING AFB DC 20332	1
EXECUTIVE OFFICER NAVAL MEDICAL RESEARCH INSTITUTE NAVAL MEDICINE COMMAND NATIONAL CAPITAL REGION BETHESDA MD 20814	1	COMMANDER US ARMY CHEMICAL BIOLOGICAL DEFENSE AGENCY ATTN AMSCB CI ABERDEEN PROVING GROUND MD 21010-5423	1
USAF ARMSTRONG LABORATORY/CFTO SUSTAINED OPERATIONS BRANCH BROOKS AFB TX 78235-5000	1	LTC RICHARD R. STOTTS BATTELLE MEMORIAL INSTITUTE JM 3 505 KING AVENUE COL UMBUS OH 43201-2695	1
DEPARTMENT OF HEALTH AND HUMAN SERVICES NATIONAL INSTITUTES OF HEALTH THE NATIONAL LIBRARY OF MEDICINE SERIAL RECORDS SECTION 8600 ROCKVILLE PIKE BETHESDA MD 20894	1	COMMANDER US ARMY MEDICAL RESEARCH INSTITUTE OF CHEMICAL DEFENSE 3100 RICKETTS POINT ROAD ATTN MCMR UV ZA MCMR UV ZB MCMR UV ZS MCMR UV RC (5 copies) MCMR UV R (11 copies) MCMR UV AI W MCMR UV D MCMR UV P MCMR UV C ABERDEEN PROVING GROUND MD 21010-5425	23
STEMSON LIBRARY ACADEMY OF HEALTH SCIENCES BUILDING 2840 RM 106 FORT SAM HOUSTON TX 78234-6100	1		
US ARMY RESEARCH OFFICE ATTN CHEMICAL AND BIOLOGICAL SCIENCES DIVISION PO BOX 12211 RESEARCH TRIANGLE PARK NC 27709-2211	1		